

# Strongyloides Infection in Hawaii: An Imported Case

Cedric M Yoshimoto MD, DTM&H\*

*Tropical diseases may present anywhere in the world. A case of strongyloidiasis in Hawaii, identified by a characteristic manifestation, larva currens, is described. Strongyloides infection may persist long after leaving an endemic area through the mechanism of autoinfection and, especially under circumstances of immunosuppression, may become overwhelming (the hyperinfection syndrome), with a likelihood of being fatal. Each case of Strongyloides infection should be treated aggressively in order to prevent this dangerous outcome; in addition, the parasite should be eliminated before immunosuppressive therapy is begun.*

## Introduction

*Strongyloides stercoralis*, the threadworm, is a nematode widely distributed in the tropics and subtropics, particularly in Southeast Asia, tropical South America and sub-Saharan Africa<sup>1,2</sup>. It also is present in the southeastern United States<sup>2,3</sup>. This human parasite has a rather complicated life cycle that sometimes results in autoinfection. In such instances, the adult female worm lays eggs that hatch into *rhabditiform larvae*, which in turn develop into invasive *filariform larvae* (a) either within the intestinal lumen or (b) through the perianal skin<sup>4,5</sup>. In either case, invasion by these larvae through the mucosa or the skin maintains or amplifies the infection in the absence of reexposure to *Strongyloides* from the external environment. In this way, people with the disease may present themselves for treatment far from endemic areas of infection.

Here we have a case to report from the Waianae Coast Comprehensive Health Center.

## Case Report

A 45-year-old Vietnamese man presented himself with a complaint of pruritic rash which had reoccurred almost monthly for the past 6 years. He had been treated for this several times previously with various medications, the names of which he did not recall, but with no relief. He reported occasional cough and shortness of breath but no wheezing, abdominal pain or diarrhea. He had lived in Vietnam until 1985, and subsequently had lived in Hong Kong for one year (when the rash first appeared), the Philippines for 6 months, and Kansas for 2 years before coming to Hawaii 2 years ago.

Examination of the skin revealed 3 separate areas (left lower chest, right abdomen, right buttock) of eruption characterized by one or more serpiginous, linear, non-indurated wheals surrounded by an erythematous flare (see Figure). The remainder of the examination was not noncontributory.

Microscopic examination of a stool specimen demonstrated *rhabditiform larvae* of *Strongyloides stercoralis* and hookworm ova.

The patient was given thiabendazole 20 mg/kg p.o. b.i.d. for 2 days and diphenhydramine 25 mg to 50 mg p.o. q.6 h. p.r.n. for the infection and for symptomatic relief of the pruritis respectively. Subsequently he was prescribed mebendazole 100 mg p.o. b.i.d. for 3 days to eliminate the infestation with hookworm.

During the following 4 months he had no recurrence of the rash; this represents the longest asymptomatic period since his affliction began. However, he failed repeatedly to provide a sample for follow-up stool examinations.

Eight family members also lived in Vietnam and were asked to provide stool samples for examination. Only one of them provided a single specimen, which was negative for *Strongyloides*.

## Discussion

Uncomplicated *Strongyloides* infection is characterized most frequently by diarrhea and abdominal pain; or it can be totally asymptomatic<sup>6,7,8</sup>.

In the case we present here, we presume that the patient acquired his infection in Vietnam and that it persisted through the mechanism of autoinfection.

*Strongyloides* is known to be widespread in Southeast Asia<sup>1,2</sup>. A high proportion of British troops who served there during World War II and were imprisoned by the Japanese became infected<sup>9,10,11</sup>, as did some American troops who served in the Vietnam war<sup>12,13</sup>. Southeast Asian refugees who emigrated to the United States and other countries have shown various rates of infection<sup>2,14</sup>.


Autoinfection may result in repeated episodes of linear urticarial cutaneous eruptions from migrating larvae, as seen in the present case. This manifestation has been termed *larva currens* (Latin: racing larva) to describe the celerity with which the larvae and consequent rash move and spread, as much as 10 cm per hour<sup>15</sup>. This is in contrast to the more deliberate pace of cutaneous *larva migrans*, which is usually caused by nonhuman hookworm larvae, and consists of an indurated linear track with little or no flare<sup>1</sup>. *Larva currens* has been described as occurring frequently among the Allied troops who had been prisoners of war in Southeast Asia during World War II<sup>9,10,11,16,17</sup>. The rash is considered by some to be pathognomonic of strongyloidiasis<sup>5,9,15</sup>. *Larva currens* is reported infrequently in other areas where *Strongyloides* infestation is endemic; it may be specific to the strain of parasite

\* Waianae Coast Comprehensive Health Center  
86-260 Farrington Highway  
Waianae, Hawaii 96792

DTM&H is a British degree, signifying Diploma in  
Tropical Medicine and Hygiene.

Submitted for publication November 5, 1991.

(Continued on page 61) ➤



**"This is my  
entire checkbook."**

Clifford Kawano, Branch Manager

## Introducing CHECK CARD.<sup>SM</sup> It makes life simpler!

Our new CHECK CARD is as easy to use as a credit card, but your purchases are deducted directly from your checking account. It's that simple! There's never any debt to worry about, and no interest charges. It's accepted everywhere VISA is, all over the world. There are no hassles, and no i.d.'s to show. Even ATM's worldwide participating in the PLUS system or VISA network accept CHECK CARD. Stop by any Central Pacific Bank branch today and we'll help you get your own CHECK CARD. All you need is a CPB checking account.



**Central Pacific Bank**  
We Put You First! Member FDIC

Main Branch / 220 S. King Street / Phone 544-0500

prevalent in Southeast Asia<sup>17</sup>.

Another consequence of autoinfection is that successive generations of parasites may sustain an infection for an extraordinary span of time. The author saw a patient in London, a former prisoner of war, who had *larva currens* for a period of about 45 years since his repatriation from Southeast Asia. Leighton and MacSween have reported a woman who suffered symptoms of strongyloidiasis for approximately 65 years<sup>18</sup>.

A third possible sequelum of autoinfection is disseminated strongyloidiasis, a so-called hyperinfection<sup>7,19,20</sup>, which may occur in both immunocompetent and immunocompromised hosts. *Filariform larvae* may invade virtually every tissue, sometimes in overwhelming numbers, leading to complications such as bowel necrosis and perforation, pulmonary consolidation, respiratory distress, and bacterial infections (sepsis and meningitis); the latter condition presumably the result of intestinal bacteria accompanying the migrating larvae<sup>5,6,20</sup>.

The case fatality rate for reported hyperinfection syndrome (HS) is very high, about 77% in immunocompromised hosts according to one review of the English language literature<sup>20</sup>. Conditions associated with HS include malnutrition<sup>7</sup>, malignancy, particularly of the lymphatic system<sup>20,21</sup>, and with treatment of corticosteroids or other immunosuppressive drugs<sup>20,22</sup>. HS has been in patients with the acquired immunodeficiency syndrome<sup>23,24</sup>, although not to a marked degree<sup>25</sup>. Of note, the urticaria and pruritis of *larva currens* may prompt treatment with corticosteroids<sup>8,18</sup>, a potentially fatal error.


The diagnosis of parasitic infections is best accomplished by finding the parasite. Stool examination, as in this case, may reveal *Strongyloides larvae*<sup>26</sup>. However, larval passage in stools is variable<sup>6</sup> and multiple examinations may be required<sup>9</sup>. Examination of duodenal contents either by aspiration<sup>27</sup> or by sampling with a retrievable swallowed nylon string (Entero-Test, HDC Corp., Mountain View, California) has been advocated<sup>28</sup>, as has sputum examination for larvae<sup>5</sup>. Pelletier and colleagues have evaluated an ELISA for antibody to *Strongyloides* larval antigen and found it useful<sup>17</sup>.

Treatment of uncomplicated strongyloidiasis involves thiabendazole<sup>29,30</sup>, albendazole, or ivermectin<sup>30,31</sup>.

In view of the possibility of subsequent HS, every patient in whom *Strongyloides* infection is diagnosed should be treated. Furthermore, every patient in whom immunosuppressive therapy is contemplated, and who may have been exposed to *Strongyloides* (eg residence in an endemic area), should be evaluated for the parasite and treated if infected<sup>2,5,6,9,20,21,22</sup>.

### Conclusions

Whether the world is growing smaller (improved communications and transportation, increased travel and migration) or becoming more tropical (global warming), tropical diseases are now a matter of potential concern in every part of the world<sup>32</sup>. It is being recognized more and more that physicians need to become familiar with this branch of medicine<sup>33,34,35,36</sup>. For example, the United States, including Hawaii, has recently experienced cases of cholera<sup>37,38</sup>, malaria<sup>39,40</sup> and dengue<sup>41</sup> reported from several states. Hawaii is the most nearly tropical of the states and it is a crossroad of Pacific travel. The medical community can expect to encounter a variety of imported, as well as indigenous exotic disorders, and should remain alert to them.



*Larva currens*: eruption on left lower chest, with several serpiginous linear wheals and surrounding erythematous flare.

### REFERENCES

1. Manson-Bahr PEC, Bell DR. *Manson's tropical diseases*. 19th ed. London, England: Bailliere Tindall, 1987.
2. Genta RM. Global prevalence of strongyloidiasis: critical review with epidemiologic insights into the prevention of disseminated disease. *Rev Infect Dis*. 1989;11(5):755-67.
3. Walzer PD, Milder JE, Banwell JG, Kilgore G, Klein M, Parker R. Epidemiologic features of *Strongyloides stercoralis* infection in an endemic area of the United States. *Am J Trop Med Hyg*. 1982;31(2):313-9.
4. Brown HW, Neva FA. *Basic Clinical Parasitology*. Norwalk, Connecticut: Appleton-Century-Crofts, 1983.
5. Grove DI. Strongyloidiasis. In: Warren KS, Mahmoud AAF, editors. *Tropical and geographical medicine*. 2nd ed. New York, NY: McGraw-Hill, 1990:393-9.
6. Beaver PC, Jung RC, Cupp EW. *Clinical Parasitology*. 9th ed. Philadelphia, Pa: Lea & Febiger, 1984.
7. Bras G, Richards RC, Irvine RA, Milner PFA, Ragbeer MMS. Infection with *Strongyloides stercoralis* in Jamaica. *Lancet*. 1964;2(7372):1257-60.
8. Milder JE, Walzer PD, Kilgore G, Rutherford I, Klein M. Clinical features of *Strongyloides stercoralis* infection in an endemic area of the United States. *Gastroenterology*. 1981;80(6):1481-8.
9. Gill GV, Bell DR. *Strongyloides stercoralis* infection in former Far East prisoners of war. *BMJ*. 1979;2(6190):572-4.
10. Gill GV, Bell DR. Longstanding tropical infection among former war prisoners of the Japanese [letter]. *Lancet*. 1982;1(8278):958-9.
11. Hill JA. Strongyloidiasis in ex-Far East prisoners of war. *BMJ*. 1988;296(6624):753.
12. Pelletier LL, Gabre-Kidan T. Chronic strongyloidiasis in Vietnam veterans. *Am J Med*. 1985;78(1):139-40.
13. Genta RM, Weesner R, Douce RW, Huitger-O'Connor T, Walzer PD. Strongyloidiasis in U.S. veterans of the Vietnam and other wars. *JAMA*. 1987;258(1):49-52.
14. Hoffman SL, Barrett-Connor E, Norcross W, Nguyen D. Intestinal parasites in Indochinese immigrants. *Am J Trop Med Hyg*. 1981;30(2):340-3.
15. Arthur RP, Shelley WB. *Larva currens*: a distinctive variant of cutaneous larva migrans due to *Strongyloides stercoralis*. *Arch Dermatol*. 1958;78(2):186-90.
16. Caplan JP. Creeping eruption and intestinal strongyloidiasis. *BMJ*. 1949;1:396.
17. Pelletier LL, Baker CB, Gam AA, Nutman TB, Neva FA. Diagnosis and evaluation of treatment of chronic strongyloidiasis in ex-prisoners of war. *J Infect Dis*. 1988;157(3):573-6.
18. Leighton PM, MacSween HM. *Strongyloides stercoralis*: the cause of an urticarial-like eruption of 65 years' duration. *Arch Intern Med*. 1990 Aug;150(8):1747-8.

(Continued on page 76) ►

# YOCON<sup>®</sup>

## YOHIMBINE HCl

**Description:** Yohimbine is a 3a-15a-20B-17a-hydroxy Yohimbine-16a-carboxylic acid methyl ester. The alkaloid is found in Rubaceae and related trees. Also in *Rauwolfia Serpentina* (L) Benth. Yohimbine is an indolalkylamine alkaloid with chemical similarity to reserpine. It is a crystalline powder, odorless. Each compressed tablet contains (1/12 gr.) 5.4 mg of Yohimbine Hydrochloride.

**Action:** Yohimbine blocks presynaptic alpha-2 adrenergic receptors. Its action on peripheral blood vessels resembles that of reserpine, though it is weaker and of short duration. Yohimbine's peripheral autonomic nervous system effect is to increase parasympathetic (cholinergic) and decrease sympathetic (adrenergic) activity. It is to be noted that in male sexual performance, erection is linked to cholinergic activity and to alpha-2 adrenergic blockade which may theoretically result in increased penile inflow, decreased penile outflow or both.

Yohimbine exerts a stimulating action on the mood and may increase anxiety. Such actions have not been adequately studied or related to dosage although they appear to require high doses of the drug. Yohimbine has a mild anti-diuretic action, probably via stimulation of hypothalamic centers and release of posterior pituitary hormone.

Reportedly, Yohimbine exerts no significant influence on cardiac stimulation and other effects mediated by B-adrenergic receptors, its effect on blood pressure, if any, would be to lower it; however no adequate studies are at hand to quantitate this effect in terms of Yohimbine dosage.

**Indications:** Yocon<sup>®</sup> is indicated as a sympatholytic and mydriatic. It may have activity as an aphrodisiac.

**Contraindications:** Renal diseases, and patient's sensitive to the drug. In view of the limited and inadequate information at hand, no precise tabulation can be offered of additional contraindications.

**Warning:** Generally, this drug is not proposed for use in females and certainly must not be used during pregnancy. Neither is this drug proposed for use in pediatric, geriatric or cardio-renal patients with gastric or duodenal ulcer history. Nor should it be used in conjunction with mood-modifying drugs such as antidepressants, or in psychiatric patients in general.

**Adverse Reactions:** Yohimbine readily penetrates the (CNS) and produces a complex pattern of responses in lower doses than required to produce peripheral a-adrenergic blockade. These include, anti-diuresis, a general picture of central excitation including elevation of blood pressure and heart rate, increased motor activity, irritability and tremor. Sweating, nausea and vomiting are common after parenteral administration of the drug.<sup>1,2</sup> Also dizziness, headache, skin flushing reported when used orally.<sup>1,3</sup>

**Dosage and Administration:** Experimental dosage reported in treatment of erectile impotence.<sup>1,3,4</sup> 1 tablet (5.4 mg) 3 times a day, to adult males taken orally. Occasional side effects reported with this dosage are nausea, dizziness or nervousness. In the event of side effects dosage to be reduced to 1/2 tablet 3 times a day, followed by gradual increases to 1 tablet 3 times a day. Reported therapy not more than 10 weeks.<sup>3</sup>

**How Supplied:** Oral tablets of Yocon<sup>®</sup> 1/12 gr. 5.4 mg in bottles of 100's NDC 53159-001-01 and 1000's NDC 53159-001-10.

#### References:

1. A. Morales et al., New England Journal of Medicine: 1221, November 12, 1981.
2. Goodman, Gilman — The Pharmacological basis of Therapeutics 6th ed., p. 176-188. McMillan December Rev. 1/85.
3. Weekly Urological Clinical letter, 27-2, July 4, 1983.
4. A. Morales et al., The Journal of Urology 128: 45-47, 1982.

Rev. 1/85



AVAILABLE AT PHARMACIES NATIONWIDE

**PALISADES  
PHARMACEUTICALS, INC.**

219 County Road  
Tenafly, New Jersey 07670  
(201) 569-8502  
1-800-237-9083

## STRONGYLOIDES INFECTION IN HAWAII

(Continued from page 61)

19. Rivera E, Maldonado N, Velez-Garcia E, Grillo AJ, Malaret G. Hyperinfection syndrome with *Strongyloides stercoralis*. *Ann Intern Med*. 1970 Feb;72(2):199-204.
20. Igra-Siegmán Y, Kapila R, Sen P, Kaminski ZC, Louria DB. Syndrome of hyperinfection with *Strongyloides stercoralis*. *Rev Infect Dis*. 1981;3(3):397-407.
21. Rogers WA, Nelson B. Strongyloidiasis and malignant lymphoma: "opportunistic infection" by a nematode. *JAMA*. 1966;195 (8):685-7.
22. Cruz T, Reboucas G, Rocha H. Fatal strongyloidiasis in patients receiving corticosteroids. *N Engl J Med*. 1966;275(20): 1093-6.
23. Piot P, Quinn TC, Taelman H, Feinsod FM, Minlangu KB, Wobin O, et al. Acquired immunodeficiency syndrome in a heterosexual population in Zaire. *Lancet*. 1984;2(8394):65-9.
24. Maayan S, Wormser GP, Widerhorn J, Sy ER, Kim YH, Ernst JA. *Strongyloides stercoralis* hyperinfection in a patient with the acquired immune deficiency syndrome. *Am J Med*. 1987; 83(5):945-8.
25. Lucas S. Missing infections in AIDS. *Trans R Soc Trop Med Hyg*. 1990;84 Suppl 1:34-8.
26. Bailey W. Laboratory diagnosis of parasitic disease. In: Manson-Bahr PEC, Bell DR, editors. *Manson's tropical diseases*. 19th ed. London: Bailliere Tindall, 1987;1489-1500.
27. Goka AKJ, Rolston DDK, Mathan VI, Farthing MJG. Diagnosis of *Strongyloides* and hookworm infections: comparison of faecal and duodenal fluid microscopy. *Trans R Soc Trop Med Hyg*. 1990;84(6):829-31.
28. Beal CB, Viens P, Grant RGL, Hughes JM. A new technique for sampling duodenal contents: demonstration of upper smallbowel pathogens. *Am J Trop Med Hyg*. 1970 Feb;19(2):349-52.
29. Grove DI. Treatment of strongyloidiasis with thiabendazole: an analysis of toxicity and effectiveness. *Trans R Soc Trop Med Hyg*. 1982;76(1):114-8.
30. Drugs for parasitic infections. *Med Lett Drugs Ther*. 1990;32(814):23-32.
31. Naquira C, Jimenez G, Guerra JG, Bernal R, Nalin DR, Neu D, et al. Ivermectin for human strongyloidiasis and other intestinal helminths. *Am J Trop Med Hyg*. 1989;40(3):304-9.
32. Royal L, McCoubrey I. International spread of disease by air travel. *Am Fam Physician*. 1989;40(5):129-36.
33. Wolfe MS. Diseases of travelers. *Clin Symp*. 1984;36(2):1-32.
34. Jong EC. Travel-related infections: prevention and treatment. *Hosp Pract*. 1989 Nov 15;24(11):145-8, 151-2, 154, passim.
35. Warrell DA. Introduction. In: Warren KS, Mahmoud AAF, editors. *Tropical and geographical medicine*. 2nd ed. New York: McGrawHill, 1990:2-4.
36. Benenson AS. Preface. In: Benenson AS, editor. *Control of communicable diseases in man*. 15th ed. Washington, D.C.:1990:xix-xxvii.
37. Centers for Disease Control. Cholera - New Jersey and Florida. *MMWR*. 1991 May 3;40(17):287-9.
38. Centers for Disease Control. Cholera - New York, 1991. *MMWR*. 1991;40(30):516-8.
39. Centers for Disease Control. Imported malaria associated with malariatherapy of Lyme disease - New Jersey. *MMWR*. 1990;39(48):873-5.
40. Centers for Disease Control. Mosquito-transmitted malaria California and Florida, 1990. *MMWR*. 1991;40(6):106-8.
41. Centers for Disease Control. Imported dengue - United States, 1990. *MMWR*. 1991;40(30):519-20.



"I've told you time and again, Harriet,  
I do NOT want a cuckoo clock for my office."